

Laser-Produced Spectra and QED Effects for Fe-, Co-, Cu-, and Zn-like Ions of Au, Pb, Bi, Th, and U

J. F. Seely, J. O. Ekberg,^(a) C. M. Brown, and U. Feldman

E. O. Hulbert Center for Space Research, Naval Research Laboratory, Washington, D.C. 20375

W. E. Behring

Laboratory for Solar Physics and Astrophysics, Goddard Space Flight Center, Greenbelt, Maryland 20771

Joseph Reader

National Bureau of Standards, Gaithersburg, Maryland 20899

and

M. C. Richardson

Laboratory for Laser Energetics, University of Rochester, Rochester, New York 14623

(Received 12 August 1986)

Spectra of very highly charged ions of Au, Pb, Bi, Th, and U have been observed in laser-produced plasmas generated by the OMEGA laser. Line identifications in the region 9–110 Å were made for ions in the Fe, Co, Cu, and Zn isoelectronic sequences. Comparison of the measured wavelengths of the Cu-like ions with values calculated with and without QED corrections shows that the inclusion of QED corrections greatly improves the accuracy of the calculated 4s-4p wavelengths. However, significant differences between the observed and calculated values remain.

PACS numbers: 32.30.-r, 52.50.Jm

The spectra of very highly charged ions are of great interest for the development of laser fusion^{1,2} and for tests of relativistic calculations of the atomic structure of

high- Z ions. In this Letter we report spectra observations of highly charged ions of Au, Pb, Bi, Th, and U ($Z = 79-92$) belonging to the FeI, CoI, CuI, and ZnI isoelectronic sequences. Comparison of the measured wavelengths with relativistically calculated theoretical wavelengths shows the importance of QED corrections in the theoretical calculations. This extends to higher Z recent work³⁻⁷ on these sequences for elements up to W ($Z = 74$).

The spectra were generated in laser-produced plasmas at the University of Rochester's Laboratory for Laser Energetics. Solid targets were spherically irradiated by 24 beams of frequency-tripled radiation (wavelength 351 nm) from the OMEGA laser.^{8,9} Each beam had an en-




FIG. 1. Comparison of the observed energies of 4s-4p transitions of Cu-like ions with energies calculated by use of Grant's program without QED corrections (open circles) and with QED corrections (solid circles).




FIG. 2. Comparison of the observed energies of the $3p^6 3d^9 2D_{5/2}-3p^5 3d^{10} 2P_{3/2}$ transitions in Co-like ions with energies calculated with Grant's program.

TABLE I. Observed and calculated wavelengths in angstroms for transitions in highly charged ions of Au, Pb, and Bi.

Sequence	Transition	Au ⁴⁹⁺ - Au ⁵³⁺				Pb ⁵²⁺ - Pb ⁵⁶⁺				Bi ⁵³⁺ - Bi ⁵⁷⁺		
		Int.	Obs.	MCDF ^a	MP ^b	Int.	Obs.	MCDF ^a	RHF ^c	Int.	Obs.	MCDF ^a
Fe I	3p ⁶ 3d ⁸ 3F ₄ - 3p ⁵ 3d ⁹ (3/2,5/2) ₃	8 ^d	24.200	23.996		2	22.420	22.250		6	21.865	21.699
Co I	3p ⁶ 3d ⁹ 2D _{5/2} - 3p ⁵ 3d ¹⁰ 2P _{3/2}	10	24.338	24.186		2	22.557	22.425		8	21.997	21.871
Cu I	4s 2S _{1/2} - 4p 2P _{1/2}	3	112.485	111.53	112.80	1	104.94	103.90	103.18	1	102.55	101.52
	4s 2S _{1/2} - 4p 2P _{3/2}	20	48.928	48.771	48.919	3	42.349	42.216	42.093	10	40.394	40.241
	4p 2P _{1/2} - 4d 2D _{3/2}	16	39.922	39.903	39.928	9	35.181	35.201	35.207	15	33.775	33.759
	4p 2P _{3/2} - 4d 2D _{5/2}	4	62.653	62.633	62.712	5	57.751	57.769	57.783	8	56.272	56.248
	4d 2D _{3/2} - 4f 2F _{5/2}	4	59.428	59.483	59.007	1	55.023	55.074	55.081	4	53.624	53.697
	4d 2D _{5/2} - 4f 2F _{7/2}	2	66.500	66.574	66.063	2	62.361	62.464	62.473	4	61.086	61.183
	4s 2S _{1/2} - 5p 2P _{3/2}	2	9.539	9.576	9.607							
	4s 2S _{1/2} - 5p 2P _{1/2}	1	10.075	10.103	10.133							
	4p 2P _{1/2} - 5d 2D _{3/2}	1	9.764	9.810	9.851							
	4p 2P _{3/2} - 5d 2D _{5/2}	4	10.887	10.919	10.966	1	9.814	9.852	9.853	7	9.480	9.530
	4d 2D _{3/2} - 5f 2F _{5/2}	5	11.804	11.840	11.901	4	10.619	10.646	10.646	2	10.253	10.287
	4d 2D _{5/2} - 5f 2F _{7/2}	5	12.100	12.137	12.210	5	10.889	10.937	10.938	3	10.542	10.576
4f 2F _{5/2} - 5g 2G _{7/2}	9	13.984	14.012		3	12.484	12.525	12.527	3	12.055	12.082	
4f 2F _{7/2} - 5g 2G _{9/2}	10	14.087	14.110		4	12.577	12.623	12.626	3	12.155	12.180	
Zn I	4s ² 1S ₀ - 4s4p 1P ₁	9	48.046	47.865 ^e		1	41.689	41.546		8	39.798	39.635

^aPresent calculation using Grant's program, Refs. 11 and 12.

^bModel potential calculation of Ivanov *et al.*, Ref. 13.

^cRelativistic Hartree-Fock calculation of Cheng and Kim, Ref. 14.

^dBlended line.

^eCalculated wavelengths of Shorer and Dalgarno, Ref. 15, are 47.676 Å (relativistic random-phase approximation) and 47.752 Å (Dirac-Hartree-Fock).

ergy of about 60 J and a duration of about 450 psec. The intensity of the focused laser beams varied from 1×10^{15} to 5×10^{16} W/cm². The spectra were photographed in the region 9–112 Å with a 3-m grazing-incidence spectrograph¹⁰ on Kodak 101-05 plates. The spectrograph was fitted with a gold-coated grating ruled with 1200 lines/mm. Light from the target was focused astigmatically onto the entrance slit of the spectrograph by a gold-coated cylindrical mirror. One laser shot was used for each spectrum.

The spectra of the high-Z ions were measured relative to spectra of O⁷⁺, Si¹⁰⁺, Si¹¹⁺, and Si¹²⁺ originating in the glass stalk and silicone adhesive used to hold the target as well as separate calibration spectra from glass microballoons. The measured wavelengths are given in Tables I and II. The uncertainty of the wavelengths is ± 0.015 Å. The intensities are visual estimates of plate darkening. The line identifications were made by isoelectronic extrapolation from lower-Z elements and by comparison with calculated wavelengths.

In Tables I and II the observed wavelengths are com-

pared with various theoretical values. The multiconfiguration Dirac-Fock (MCDF) values with transverse Breit and QED corrections were calculated by us with the computer program of Grant *et al.*¹¹ and McKenzie, Grant, and Norrington.¹² The QED calculations in this program include second-order vacuum polarization corrections as prescribed by Fullerton and Rinker¹⁷ and self-energy corrections obtained by scaling the hydrogenic $n=1,2$ results of Mohr^{18,19} according to $1/n^3$.

For the Cu-like ions the QED corrections have a measurable effect only for the 4s-4p transitions. In Fig. 1 we compare the observed wavelengths for 4s-4p transitions in Cu-like ions with wavelengths calculated with the use of Grant's code, both with and without the QED corrections. These calculations were made in the extended average level mode. The energy functional optimized was the statistically weighted average of the energies of the two fine-structure levels at each nl term. The differences are seen to vary systematically. Clearly the inclusion of QED corrections in the calculations improves the accuracy of the results. However, significant differ-

TABLE II. Observed and calculated wavelengths in angstroms for transitions in highly charged ions of Th and U.

Sequence	Transition	Th ⁶⁰⁺ -Th ⁶³⁺			U ⁶²⁺ -U ⁶⁵⁺		
		Int.	Obs.	MCDF ^a	Int.	Obs.	MCDF ^a
Co I	$3p^6 3d^9 2D_{5/2} - 3p^5 3d^{10} 2P_{3/2}$				3	17.561	17.483
Cu I	$4s^2 S_{1/2} - 4p^2 P_{3/2}$	5	28.990	28.839	2	26.400	26.235
	$4p^2 P_{1/2} - 4d^2 D_{3/2}$	3	25.180	25.170			
	$4p^2 P_{3/2} - 4d^2 D_{5/2}$	3	46.830	46.816			
	$4d^2 D_{5/2} - 4f^2 F_{7/2}$	2	53.154	53.255			
Zn I	$4s^2 1S_0 - 4s 4p^1 P_1$	2	28.703	28.547	1	26.175	26.000 ^b

^aPresent calculation by use of the program of Grant *et al.*, Ref. 11, and McKenzie, Grant, and Norrington, Ref. 12.

^bCalculated values of Shorer, Ref. 16, are 25.847 Å (relativistic random-phase approximation) and 25.876 Å (Dirac-Hartree-Fock).

ences between observed and calculated values still remain.

Figure 2 shows the differences between the observed wavelengths of the $3p^6 3d^9 2D_{5/2} - 3p^5 3d^{10} 2P_{3/2}$ transitions in Co-like ions and those calculated with the Grant code. Except for the transition in Yb⁴³⁺ ($Z=70$), which was considered to be poorly determined,⁶ the present measurements are consistent with those of the lower- Z ions.⁶

In conclusion, the agreement between the presently measured transition energies and the transition energies calculated with the Grant code improves with increasing Z except for the Cu-like $4s^2 S_{1/2} - 4p^2 P_{1/2}$ and $4s^2 S_{1/2} - 4p^2 P_{3/2}$ transitions and the Zn-like $4s^2 1S_0 - 4s 4p^1 P_1$ transition. For these transitions, the discrepancies grow with Z and indicate a departure from the scaled hydrogenic QED contribution to the $4s$ energy.

We thank W. J. Powers and A. E. Williams for providing some of the target materials. The experiments were coordinated by F. J. Marshall. We thank the target fabrication group and the laser operations staff at the Laboratory for Laser Energetics for their expert technical assistance. This work was supported by the U.S. Department of Energy.

^(a)Also at Sachs/Freeman Associates, Landover, MD 20785. Permanent address: Department of Physics, University of Lund, Lund, Sweden.

¹W. C. Mead *et al.*, Phys. Fluids **26**, 2316 (1983).

²D. L. Matthews *et al.*, J. Appl. Phys. **54**, 4260 (1983).

³J. Reader and G. Luther, Phys. Rev. Lett. **45**, 609 (1980).

⁴J. Reader and G. Luther, Phys. Scr. **24**, 732 (1981).

⁵J. Reader and A. Ryabtsev, J. Opt. Soc. Am. **71**, 231 (1981).

⁶J. Reader, J. Opt. Soc. Am. **73**, 63 (1983).

⁷N. Acquista and J. Reader, J. Opt. Soc. Am. B **1**, 649 (1984).

⁸J. M. Soares, R. J. Hutchinson, S. D. Jacobs, L. D. Lund, R. L. McCrory, and M. C. Richardson, in *Proceedings of the Tenth Symposium on Fusion Energy, Philadelphia, Pennsylvania, 1983* (IEEE, New York, 1983), p. 1392.

⁹M. C. Richardson *et al.*, in *Proceedings of the Seventh Meeting on Laser Interaction and Related Plasma Phenomena*, edited by H. Hora and G. H. Miley (Plenum, New York, 1986), Vol. 7.

¹⁰W. E. Behring, R. J. Ugiansky, and U. Feldman, Appl. Optics **12**, 528 (1973).

¹¹I. P. Grant, B. J. McKenzie, P. H. Norrington, D. F. Mayers, and N. C. Pyper, Comput. Phys. Commun. **21**, 207 (1980).

¹²B. J. McKenzie, I. P. Grant, and P. H. Norrington, Comput. Phys. Commun. **21**, 233 (1980).

¹³L. N. Ivanov, E. P. Ivanova, E. Ya. Kononov, S. S. Churilov, and M. A. Tsirekidze, Phys. Scr. **33**, 401 (1986).

¹⁴K.-T. Cheng and Y.-K. Kim, At. Data Nucl. Data Tables **22**, 547 (1978).

¹⁵P. Shorer and A. Dalgarno, Phys. Rev. A **16**, 1502 (1977).

¹⁶P. Shorer, Phys. Rev. A **18**, 1060 (1978).

¹⁷L. W. Fullerton and G. A. Rinker, Phys. Rev. A **13**, 1283 (1976).

¹⁸P. J. Mohr, Ann. Phys. **88**, 52 (1974).

¹⁹P. J. Mohr, Phys. Rev. Lett. **34**, 1050 (1975).